



sea state
cci


User Requirements Document (URD)

version 2.0, 14 October 2021

Contents

1. Introduction	4
1.1 Purpose	4
1.2 Scope	5
1.3 Description of the main sea state variables	5
1.4 Document structure	6
2. Review of previous user requirements	7
2.1 Sea states as part of the Earth system	7
2.2 EUMETSAT Observation Requirements for Nowcasting in 2015-2025	7
2.2.1 Observational performance level	8
2.2.2 Priority of the requirements	9
2.3 Requirement defined by GCOS-200 (2016)	10
2.4 Proposition for updating requirements (see [15])	11
3. Application areas for sea state climate data	12
3.1 Climate research and engineering applications	12
3.2 Modelling activities	13
3.3 Remote sensing	13
3.4 Air-sea fluxes	13
3.5 Other wave-related effects in the Earth System	14
4. User surveys	15
4.1 Introduction	15
4.2 First user survey	15
4.2.1 Users involved in the survey	15
4.2.2 Field of applications	16
4.2.3 The importance of various sea state parameters	17
4.2.4 Temporal coverage and spatial resolution	18
4.2.5 Interest in sea state data	18
4.2.6 Interest in other Essential Climate Variables	19
4.2.7 Participation to the User Consultation Meeting and training session	19
4.3 Second user survey	20
4.3.1 User profile and study area	20
4.3.2 Use of sea state data	23
4.3.3 Experience with satellite data	25
4.3.4 Experience with Sea State CCI products	25
5. Consolidated list of User Requirements	29
5.1 Top level requirements	29
5.2 Specific data requirements	30
6. References	33
Appendix 1 - Open comments from survey participants	37

Authors	Approved	Signature	Date
Fabrice Ardhuin and Guillaume Dodet	Fabrice Ardhuin, Ellis Ash		14/10/2021
ESA Approval			

Issue	Date	Comments
1.0	06/02/2019	First complete version for review by ESA
2.0	14/10/2021	Updates for final project year including results of a new requirements questionnaire from UCM2

1. Introduction

1.1 Purpose

The purpose of the *User Requirements Document* is to provide a complete set of individual requirements and constraints to meet the needs of the Climate Research Community for ECV data products related to sea state. We also note that the definition of the present wave climate may be just as important at the detection of its evolution. More generally, there are many users of the same wave data that do not belong to the climate research community, and their needs are considered here too.

As stated by GCOS (http://tiny.cc/GCOS_seastate) “*Sea state is best known for its impacts on marine safety, marine transport and damage to structures. However, waves also affect the growth or decay of sea ice, beach erosion, surface albedo, gas transfer, transport of larvae and contaminants such as oil, and air–sea exchange of energy, moisture and momentum. They thereby play large roles in the global cycles of energy, water and carbon.*” Routine observation of sea states by satellites go back 25 years, and satellite data has been key in defining and adjusting parameterizations in numerical wave models. Altimeters, SARs, high resolution optical imagery and the new SWIM wave scatterometer on CFOSAT have been developed and successfully utilized to measure significant wave height (hereinafter H_s), mean square slope (mss), and swell partitions information (height, period and direction). In climate change studies based on satellite data, it is a major challenge to construct homogeneous time series from a series of consecutive satellite sensors needed for detection of changes over several decades.

At the same time there is an evolution in sensors and observation technology, which makes it possible to measure new sea state parameters for future monitoring, for example foam properties using passive radiometers such as SMOS. Also a number of other sea state parameters are important for climate research, such as the sea state bias in altimeter data or the directional slope variance tensor for sunglint in ocean color images. There is thus a general need for a consistent estimate of all these variables.

The present document, guided by user input, is strongly focused on usual sea state variables and their long term variability and evolution, still keeping an eye on the other uses of wave data. The biggest headlines about sea states have been associated with publications on observed trends (Young et al. 2011) that do not always agree with modeled trends in some regions such as in the North Pacific (e.g. Semedo et al., 2010), although this is particularly complex where tropical storms dominate the extremes. One of the goals in wave climate research is to better understand the natural variability of sea states and better define the statistics of the extremes (which are so important for engineering design application and natural hazard mitigation) in a context of global change with positive and negative drifts. And even where the wave height trends may be negative, defining uncertainties is critical for the understanding of total sea level changes.

1.2 Scope

The scope of this URD is to identify, analyze and assess the requirements for sea state data specified by the sea state user community, with a focus on the modelling and climate research community. This URD is based on literature review, two dedicated user surveys, and the analysis of the requirements by the Climate Research Group within the consortium and representatives from the Climate Modelling User Group (CMUG).

The requirements will incorporate the relevant sea state parameters needed by the research community, with focus on product, accuracy, coverage, resolution and stability of the measurements, error characterisation, quality flags, metadata (including processing algorithms), product formats, grid. The requirements for validation data from non-space platforms are also addressed. The requirements for sea state ECV data include several sea state parameters in addition to the Hs and spectral partition products.

The requirements are defined for a number of applications in particular climate modelling including model development, model validation, model initialisation, boundary layer definition and data assimilation, and climate research based on time series analysis. User requirements from other application areas such as operational marine weather forecasting, ocean engineering, marine transportation and offshore operations, remote sensing and seismology are also included. The requirements are divided in two categories: (1) minimum requirements to sea state data in order to be useful for the different applications (“must have”), and (2) target requirements, which are expected to be obtained by EO data after careful validation and/or merging with non-EO data (“nice to have”). The requirements will also account for foreseen needs as sea state and climate models become further developed.

1.3 Description of the main sea state variables

Significant wave height (Hs) is the most common sea state parameter and is defined as 4 times the standard deviation of the surface elevation in a record that is typically 20 minutes long. Hs is used in all applications, from navigation safety to coastal engineering. The acronym **SWH** is also used for that same quantity in most remote sensing publications.

Besides Hs, the time and/or spatial scales of the waves are of interest because they determine the energy flux associated to the wave field (proportional to the wave period for linear waves), forces on structures, the extent of the coastal run-up, among others. Also, the direction of wave propagation is relevant for many aspects including the impact on shorelines and navigation hazards. More details can be found in Ardhuin et al. (2019).

Hence the sea state is generally described by the **directional wave spectrum $E(k, \theta)$** . Most sea state variables of interest can be derived from the spectrum. These include,

- The wavenumber spectrum $E(k) = \int_0^{2\pi} E(k, \theta) d\theta$
- frequency spectrum $E(f) = \int_0^{2\pi} E(k, \theta) dk/df d\theta$, in which dk/df is uniquely defined for linear waves by the dispersion relation.

In general this dispersion relation requires a knowledge of the wave direction and the effective current¹ velocity vector \mathbf{U}

$$(2\pi f) = [gk \tanh(kD)]^{1/2} + \mathbf{k} \cdot \mathbf{U}$$

Hence, when currents cannot be neglected, it is not straightforward to transform a frequency spectrum $E(f)$ as measured by a moored buoy, to a wavenumber spectrum $E(k)$ as measured by a space-borne radar.

Other moments of interest include

- directional moments $a_n(f) = \int_0^{2\pi} \cos(n\theta) E(k, \theta) dk/df d\theta / E(f)$ and $b_n(f) = \int_0^{2\pi} \sin(n\theta) E(k, \theta) dk/df d\theta / E(f)$
- Moment periods $T_{mp,q}$, in particular the mean period $T_{m0,2}$ and the energy period $T_e = T_{m0,-1}$ defined as $T_{mp,q} = \left[\frac{\int_0^{fmax} E(f) f^p df}{\int_0^{fmax} E(f) f^q df} \right]^{1/q}$ that can be sensitive to the maximum frequency used in the integration
- the mean square slope $m_{ss} = \int_0^{\infty} k^2 E(k) dk$
- partitions and their moments (Gerling 1992, Hanson and Phillips 2001)

Relevant directional parameters are computed from the directional moments for a particular range of frequencies or spectral partition, e.g., the mean direction and spread from the first moments at each frequency (Kuik et al. 1988)

$$[\cos(\theta_1(f)), \sin(\theta_1(f))] = [a_1(f), b_1(f)] \quad \text{and} \quad \sigma_1 = [2 - (a_1^2(f) + b_1^2(f))]^{1/2}$$

Besides \mathbf{H}_s , all other parameters are estimated indirectly from remote sensing data. We also note that the full spectrum $\mathbf{E}(\mathbf{k}, \theta)$ is generally not accessible in routine in situ measurements, only directional moments are available from which a directional spectrum may be estimated (e.g. Lygre and Krogstad 1986).

1.4 Document structure

This User Requirements Document is organised into the following sections:

- Section 2. Review of previous user requirements
- Section 3. Application areas for sea state climate data
- Section 4. User survey
- Section 5. Consolidated requirements list
- Section 6. References

¹ "Effective" in the sense that it is the velocity that advects the phase of the wave train, and is generally a function of the current profile and the wavenumber (e.g. Stewart and Joy 1974, Andrews and McIntyre 1978, Kirby and Chen 1989).

2. Review of previous user requirements

2.1 Sea states as part of the Earth system

Sea state indeed affects all activities at sea (shipping, oil & gas exploration and exploitation, fish farming) on the coast (harbours, coastal defence, marine renewable energy systems, recreational uses). Many activities require more and more accurate information from extremes (design criteria for structural failure) through to calm windows (for maintenance of offshore wind farms). Beyond activities directly linked to the ocean, sea states are of general interest in the Earth system. For example, ocean waves largely define air-sea fluxes and upper ocean mixing (Jähne and Haußecker 1998, Veron 2015, d'Asaro 2016), sediment resuspension, transport and coastal geomorphology. Waves are also the source of most of the recorded seismic noise that can be used for solid Earth monitoring (e.g. Shapiro et al. 2005) and wave climate analysis (Bernard 1990, Grevemeyer et al. 2000).

In return, extreme waves and their trace in the geological record are used as evidence for past storminess using paleo-shorelines (Bouchette et al. 2010), ripple marks (Allen et Hoffman 2005) or wave-transported boulders (Hansen et al. 2016). It is thus very important to link extreme sea states to these geological marks under present climate conditions from shoreline features (Ashton et al. 2001) to ripples (e.g. Ardhuin et al. 2002), and boulders (Kennedy & al. 2016, Autret et al. 2016, Cox 2018), in order to better understand the geological record and past climates.

Monitoring and forecasting of sea states is closely integrated with ocean and atmospheric observations and modelling. Satellite EO data play an essential role in observing atmospheric and ocean variables, including sea states.

Observations of the ocean are required for monitoring of climate and the environment at seasonal-to-interannual-to-decadal time scales. In particular, the availability of operational ocean observations is prerequisite for quality weather and ocean state forecasts. Already today, global and regional numerical weather prediction models, seasonal to inter-annual forecasts and climate models assimilate ocean observations to generate initial conditions or boundary conditions.

Requirements for sea state data are therefore closely linked to requirements for ocean and atmospheric observations and modelling, with a recent strong interest in the polar regions. Several user requirement documents have been prepared for ocean and sea ice observation from satellites. The following sections give an overview of some of these documents.

2.2 EUMETSAT Observation Requirements for Nowcasting in 2015-2025

As listed in [1] the impact of improvements to observations has been assessed for the forecast service requirements using appropriate nowcasting and very short range forecasting techniques and have then been analysed to identify the key breakthroughs. The dominant forecasting method in 2015-2025 is expected to be Numerical Weather Prediction (NWP) which by then will be able to resolve the scales of interest in very short range forecasting.

The respective main observational requirements for applications of the above user groups are summarised in this table:

	A	B	C	D	E	F	G	H
1	Title	Sub-Title	Required variable	Conditions (1)	Applications	Nearest WMO/CEOS parameter	WMO/CEOS code	Required Altitude Range (2)
52			Ocean					
53	Observation & Extrapolation	Ocean	Sig wave ht		Marine transport	Significant wave height	72	surface
54	Observation & Extrapolation	Ocean	Sig wave ht	near shore	Marine construction	Significant wave height	72	surface
55	Observation & Extrapolation	Ocean	Wave period		Marine transport	Dominant wave period	73	surface
56	Observation & Extrapolation	Ocean	Wave direction		Marine transport	Dominant wave direction	74	surface
57	Observation & Extrapolation	Ocean	Surge height	near shore	Life - Coastal flooding	Sea level	75	surface

I	J	K	L	M	N	O	P	Q	R	S	T	U	V
Accuracy threshold (3)	Accuracy optimum (3)	dx threshold	dx optimum	dz threshold	dz optimum	dt threshold	dt optimum	delay threshold	delay optimum	Priority level	Breakthrough level (if any)	Current Observing Techniques	Level met
0.5m	0.1m	50km	1km	-	-	6hr	1hr	2hr	30min	VH	0.5m / 10km / 1hr	Surface in situ or HF radar	N
0.2m	0.1m	10km	1km	-	-	3hr	1hr	1hr	5min	H	0.2m / 10km / 1hr	Surface in situ or HF radar	N
1s	0.5s	50km	1km	-	-	6hr	1hr	2hr	30min	H	1s / 10km / 1hr	Surface in situ or HF radar	N
20°	10°	50km	1km	-	-	6hr	1hr	2hr	30min	M	20° / 10km / 1hr	Surface in situ or HF radar	N
20cm	10cm	50km	1km	-	-	6hr	1hr	2hr	30min	VH	0.2m / 10km / 1hr	Surface in situ	[U]

Table 2.1: requirements as defined by [1]

Those requirements were extensively documented before and examples of respective discussions can be found in the position papers produced by the EUMETSAT Application Expert Groups during the Meteosat Third Generation (MTG) definition process [1,2,3]. Further input has been retrieved from the IGOS (Integrated Global Observing Strategy) Ocean Theme Paper [4] and the report of the Intergovernmental Oceanographic Commission (IOC/UNESCO) on Observing the Oceans in the 21st Century [5] and the WMO and GOOS [6,7] requirements published online.

Furthermore, our requirements consider and where necessary, build upon, those contained in the following documents:

- GMES, Sentinel 3 [8]
- NPOESS IORD/II [9]
- GCOS and WCRP (via the WMO on-line requirements) [6]
- EUMETSAT OSISAF [10]

All user groups continue to require comprehensive, accurate and higher resolution oceanographic satellite observations, driven by the increase in model resolution and the number of assimilated variables.

2.2.1 Observational performance level

Three performance levels are defined as follows:

1) **Threshold** is the limit below which the observation becomes ineffectual and is of no use for the targeted application

2) **Breakthrough** level represents the level beyond which a significant improvement in the target application is achieved.

3) **Objective** is the maximum performance limit for the observation, beyond which no significant improvement in the targeted application is achieved.

Accuracy: For operational meteorological applications, the accuracy is the root mean square (r.m.s.) difference between the actual measurement and the truth, inclusive of random errors and bias. This assumes that the main source of error relevant to the 'single level 2' measurement is the random component, the bias error being small enough to not significantly influence subsequent mission definition.

Spatial resolution : Horizontal resolution, Δx - The horizontal resolution is the minimum horizontal spatial scale that must be resolved by the observing system. In most cases, the horizontal resolution is more or less transferable to the resolution of a potential instrument assuming some appropriate sampling which will be finalised during the instrument design phase.

Observation Cycle, Δt : 'Observation cycle' or 'revisit time' is the time elapsed between measurements over a given location. By default, the observation cycle is applicable to the whole Earth surface including the equatorial and polar regions, so that Δt is the time needed to cover the whole Earth surface with at least one measurement. The exception is for high latitude requirements, where the observation cycle of any polar orbiting satellite is specific for the particular latitude and is not applicable to equatorial regions.

Delay, δ

The 'delay' is the time elapsed between observation by the satellite and the availability the product to user interface, including the nominal dissemination time.

Additional Notes for Climate Requirements

Some climate requirements are given as a 'level 3' product, which is the average of a series of 'level 2' measurements taken over a time period, Δt , (or sometimes distance, Δx) in order to reduce the uncertainty and sampling variability to a sufficiently low level. The spatial and temporal resolutions then refer to the integration periods. However, despite this, the r.m.s. accuracy for each single 'level 2' measurement is given in the accuracy requirement and not the corresponding averaged product accuracy.

2.2.2 Priority of the requirements

The following codes are used to prioritise the requirements listed in Table 2.1:

Priority 1 (Very High): Mandatory requirements that drive the mission, these requirements are of utmost importance for the success of the mission and must be implemented.

Priority 2 (High): Important requirements that substantially contribute to the success of the mission. Reasonable effort shall be made to implement them.

Priority 3 (Medium): Beneficial requirement that has certain value to the success of the mission, it shall be implemented with minimum effort.

Priority 4 (Low): Requirements which are marginally contributing to the success of the mission. It shall only be implemented on an opportunistic basis. No dedicated effort will be made to implement them.

The priorities assigned in the user requirements table 2.1 are technology free and are thus independent of the availability of appropriate reliable and affordable measurement techniques.

2.3 Requirement defined by GCOS-200 (2016)

After earlier versions with much higher resolution goals, the sea state observation requirements have been defined in the most recent GCOS document (GCOS report no. 200: The Global Observing System for Climate: Implementation Needs).

These requirements are reproduced in Table 2.2.

Ocean ECV product requirements								
ECV	Products	Frequency	Resolution	Required measurement uncertainty	Stability (per decade unless otherwise specified)	Standards/References	Entity (see Part II, section 2.2) ¹⁰⁰	
							Satellite	In situ
Sea-surface temperature	Sea-surface temperature	Hourly to weekly	1–100 km	0.1 K over 100-km scales	< 0.03 K over 100-km scales	See EOVS Specification sheets www.gooscean.org/eov	WGClimate	JCOMM
Subsurface temperature	Interior temperature	Hourly to monthly	1–10 km	0.01 K	Not specified			JCOMM
Sea-surface salinity	Sea-surface salinity	Hourly to monthly	1–100 km	0.01 psu	0.001 psu		WGClimate	JCOMM
Subsurface salinity	Interior salinity	Hourly to monthly	1–10 km	0.01 psu	Not specified			JCOMM
Surface currents	Surface geostrophic current	Hourly to weekly	30 km	5 cm/s	Not specified		WGClimate	JCOMM
Subsurface currents	Interior currents	Hourly to weekly	1–10 km	0.02 m/s	Not specified			JCOMM
Sea level	Global mean sea level	Weekly to monthly	10–100 km	2–4 mm (global mean); 1 cm over a grid mesh	< 0.3 mm/yr (global mean)		WGClimate	JCOMM
	Regional sea level	Hourly to weekly	10 km	1 cm (over grid mesh of 50–100 km)	< 1 mm/yr (for grid mesh of 50–100 km)		WGClimate	JCOMM
Sea state	Wave height	3-hourly	25 km	10 cm	5 cm		WGClimate	JCOMM
Surface stress	Surface stress	Hourly-monthly	10–100 km	0.001–4 Nm ²	Not specified			JCOMM

Table 2.2. Requirements for horizontal resolution, temporal resolution, measurement accuracy and stability over a decade for the ocean ECVs as defined by GCOS.

As stated by GCOS (http://tiny.cc/GCOS_seastate) “Sea state is best known for its impacts on marine safety, marine transport and damage to structures. However, waves also affect the growth or decay of sea ice, beach erosion, surface albedo, gas transfer, transport of larvae and contaminants such as oil, and air–sea exchange of energy, moisture and momentum. They thereby play large roles in the global cycles of energy, water and carbon.”

Product	Frequency	Resolution	Required uncertainty	Required stability (per decade)
Hs	3-hourly	25 km	10 cm	5 cm

Given that Hs is mostly provided by satellite altimeters and that there are at most 6 altimeters flying at any given time, these requirements cannot be attained by existing satellite data for the period 2003-2020. Further, coastal applications generally require even finer resolution, of the order of 100 m in space (Camus et al. 2013, Boudière et al. 2014). Likewise, a higher temporal resolution of the order of 1 hour is necessary where modulation by tides are important. This is particularly relevant for extreme water levels that combine wave run up and storm surges. Today numerical models and/or statistical methods are used to arrive at these resolutions.

However, these requirements are not mutually consistent. The urgency of understanding total sea level at the coast (e.g. Melet et al., 2018) is clearly calling for a stability that matches that of the offshore sea level. Typically, 1 cm increase in offshore Hs gives 0.5 to 1 cm increase in maximum sea level at the coast (Poate et al. 2016, Dodet et al. 2018). It is not unreasonable to ask for 2 mm/year accuracy for Hs when the requirement for regional sea level is at 1 mm/year. This is particularly important in today’s transition where the total ice-shelf melt contribution to sea level rise is still limited to a few centimeters. In the long term, with sea level rise of several meters, the few centimeters to decimeters due to waves will probably be less important, except where changes are dramatic, as is the case in the Arctic (e.g. Stopa et al., 2016) and possibly in tropical cyclones (Shimura et al., 2016).

2.4 Proposition for updating requirements (see [15])

Based on the analysis of all previous requirements, the evolution of user needs, as further discussed in the next sections, and the need for consistency with other ECVs we propose to update the GCOS-200 numbers for Hs and define requirements for other variables.

	Variable	Frequency	Resolution	Uncertainty	Stability
GCOS-200	H_s	3-hourly	25 km	10 cm	5 cm/decade
GCOS-200	regional sea level	hourly	10 km	1 cm	< 1 mm/year
WMO	H_s	??	??	5-10% or 10 to 25 cm	??
WMO (2017)	$T_{m0,2}$??	??	0.1 to 1 s	??
WMO (2017)	θ_m	??	??	10 deg	??
this paper	global to regional H_s	3-hourly	25 km	10 cm or 5%	5 cm/decade
this paper	coastal H_s	1-hourly	1 km	10 cm or 5%	< 1 mm/year
this paper	regional $T_{m0,-1}$	3-hourly	25 km	0.2 s	< 0.1 s/decade
this paper	regional $T_{m0,2}$	3-hourly	25 km	0.2 s	< 0.1 s/decade

Table 2.3 : proposed updates to sea state requirements [15]. “Coastal” can be understood as regions where proximity to land and/or shallow water has a particular impact on the sea state. This thus includes all waters shallower than 100 m, or with a distance to coast shorter than 200 km.

These should be understood as objectives. In particular the objective for trends on H_s should apply to both the mean value and extreme values (up to percentile 99 and 100 year return period).

3. Application areas for sea state climate data

Sea state climate data are used in a wide range of application areas, ranging from climate research activities to marine biology and ecosystems research, management of marine resources, sea transportation, offshore exploration, design and construction of vessels and platforms, impact on indigenous people, insurance, governance and policy making. In this project focus is on climate research and modelling activities, but also requirements to other user groups are considered.

3.1 Climate research and engineering applications

Even though remote sensing data alone cannot reach these resolutions needed in coastal applications, remote sensing is critical for:

- validating and calibrating numerical models offshore (spatially coarse data is generally enough, see e.g. Stopa et al. 2016)
- validating patterns and gradients (in coastal regions, near the ice edge, over current gradients, at the peak of storms...): this has not been applied much so far because:
 - routine altimeter processes are dominated by noise for along-track wavelengths under 80 km or so (Arduin et al. 2017b). As new processing techniques have been proposed (e.g. Passaro et al. 2014, Halimi et al. 2016), there are great opportunities to better resolve these variations.
 - few images of wave parameters were available so far. This can change with more widely available SAR (e.g. Gemmrich et al. 2016, Arduin et al. 2017a) or sun glitter imagery (Kudryavtsev et al. 2017).

Finally, without even mentioning its evolution with global change, the wave climate, including extremes often defined through H_s statistics associated with long recurrence intervals (i.e. 20, 50, 100yr return period) is a key element in the design and operation of ocean and coastal infrastructures.

The impact of climate change on the wave climate and its applications is a topic of active research and is mostly limited by the poor knowledge on trends and variability of extreme sea states (e.g. Bitner-Gregersen et al. 2013).

As a result, a clear requirement is a validation of the highest values of H_s for all regions of the world ocean. Hanafin et al. (2012) used wind speeds and the periods of radiated swells as a consistency check on the 20.1 m H_s found in the North Atlantic storm Quirin. Cardone et al. (2015) chose to filter the data without any discussion on the filter properties.

Due to the particular temporal sampling of waves measured by altimeters, the statistical distribution must be corrected for sampling biases (Izaguirre et al, 2011). Figure 3.1 shows an example with the 10-year H_s return values as derived from the altimeter record, which is within 1 m of the same parameter derived from a model hindcast sampled at the satellite measurements in time and space, but differs by up to 20% from a full model time series.




Figure 3.1: (a) Estimation of a 10-year return period value of H_s using the GEV distribution from the altimeters, (b) difference between the altimeters and colocated WW3 data, (c) difference for full time series of modeled data (taken from Stopa et al. presentation at 2016 OSTST Meeting, La Rochelle).

3.2 Modelling activities

Wave model development and the assessment of the suitability of the necessary wind forcing is generally done by comparing model output to *in situ* observations, if available. However, it is seldom the case that one has enough *in situ* observations to cover the full extent of the model domain. For this reason, the wave modelling community relies extensively on the comparison of the model output with all available altimeter observations. Provided that the area covered by the model domain is large enough and that the period covered by the model simulation includes enough interesting weather, the statistical analysis of the model-altimeter along track collocations is a useful tool in development phase of the model as well as for assessment of the global characteristics of the modelling system (Wiese et al. 2018).

3.3 Remote sensing

Because ocean waves have clear signatures in most ocean remote sensing techniques, either adding noise or biases, stable corrections and detection is very important for:

- sea level estimates from altimetry (e.g. Tran et al. 2010, Passaro et al. 2018) and tide gauges (e.g. Aucan et al. 2012)
- glitter and foam contamination in ocean color imagery
- the surface mean square slope and foam cover and thickness are major sources of uncertainty in surface salinity retrieval (e.g. Reul and Chapron 2003).
- waves have an impact of retrieved wind speeds from all sensors (altimeters, scatterometers, radiometers), that is not fully understood despite recent attempts to reconcile these different records (Young and Donelan 2018).

3.4 Air-sea fluxes

Properties of the ocean and atmospheric mixed layers in which most human activity takes place, is largely driven by the air-sea exchanges of heat, water, momentum, gasses. These fluxes have been parameterized with resistance laws in which coefficients are often a function of wind speed alone. Additional dependencies on sea state properties have been strongly debated for the momentum flux (Drennan et al. 2005), while more recent analyses show a moderate impact for intermediate wind speeds (Edson et al. 2013), even though it is expected that the surface roughness caused by waves should play a role (Donelan 2018) although one that is often correlated with the wind speed.

The impact of waves on upper ocean mixing and sea surface temperature is clear, in particular in cases of shallow mixed layers (e.g. Noh et al. 2011, Janssen 2012). This has important consequences on air-sea heat fluxes (Hansen et al. 2011).

Particular uncertainties in the climate systems are associated with cloud nucleation, which relies on marine aerosol production (e.g. Veron 2015) accounting for up to 30% of cloud nucleation (Quinn et al. 2018). The source functions for marine aerosols is expected to depend on sea state parameters, such as the height of breaking waves for which the significant wave height may be a good proxy away from the swell-dominated regions (de Leeuw et al. 2011).

3.5 Other wave-related effects in the Earth System

The interactions of waves and sea ice certainly influence ice edge dynamics and ice properties near the edge (Squire et al. 1995, Stopa et al. 2018). In particular, waves are associated with the formation of pancake ice (e.g. Doble et al. 2003) that is the most common type of ice formation in the Southern ocean, and is becoming increasingly important in the Arctic too (Thomson et al. 2018). Wave action over fragmented ice can have an important influence on the ice thickness (Sutherland and Dumont 2018).

4. User surveys

4.1 Introduction

In order to develop the Essential Climate Variables (ECVs) for sea state, it is necessary to perform user surveys and analyse the requirements extracted from these surveys. During the phase 1 of the Sea State CCI project, two questionnaires were implemented in order to collect information on the user's experience with satellite wave data, the wave parameters they are mostly interested in, their requirements in terms of spatial resolution and time coverage, and their intention to use the data in relation with other ECVs. The main results of these surveys are described in the following sections and open comments from participants are listed in Appendix 1.

4.2 First user survey

The user survey was conducted from 11 December 2018 to 25 January 2019. This survey was developed using an online Google form (<https://goo.gl/forms/1GGPGc4APA1PXCax2>) and was broadcasted on several mailing lists (including globwave users, IOWAGA users and the "coastal list" managed by the University of Delaware), as well as via Twitter.

4.2.1 Users involved in the survey

As of January 25, a total of 184 participants had replied to the questionnaire. From this survey we see that the participants are mostly working in academic and research institutions (75%) spread all over the world (Figure 4.2.1, 4.2.2 and 4.2.3). The six most represented countries are France (18%), the United States (13%), Italy (8%), the United Kingdom (6%) and Australia (4%) ex-aequo with Spain (4%). The 43% remaining participants are based on the American (Canada, Mexico, Costa Rica, Colombia, Brasil, Uruguay, Chile), European (Belgium, Denmark, Germany, Norway, Ireland), Asian (China and India) and African continents (Tunisia, Egypt, Benin, Ghana, South Africa).

You: your professional situation

184 responses




Figure 4.2.1. Professional situation of the participants




Figure 4.2.2. Countries where participants currently do most of their work (in blue)




Figure 4.2.3. The 10 most represented countries in the survey

This spatial distribution of the participants highlights the worldwide interest for high quality sea state information.

4.2.2 Field of applications

The participants had to choose between eight fields of applications. The results show that the majority (66.3%) of the participants are working in the field of oceanography (38.6%) and coastal engineering (27.7%). A significant fraction of participants working in the fields of climate (8.7%) and marine meteorology (8.7%) also took part in the survey (Figure 4.2.4).

You: what is your main field of applications

184 responses




Figure 4.2.4. Distribution among the field of applications

4.2.3 The importance of various sea state parameters

The most important parameter for the quasi totality of the participants is **the significant wave height, followed by the wave period (peak period T_p and mean period T_{m02} rather than the mean period T_{m01} , not shown here) and the wave direction**. The 2-D wave spectrum $E(f,\theta)$ or $E(k,\theta)$ is also of interest for the majority of the participants, and overtakes the swell partition and the heave spectrum. Among the other altimetry-derived parameters, the mean square slope of the sea surface and the microseism sources are also relevant.




Figure 4.2.5 Sea state parameters that participants would like to use

4.2.4 Temporal coverage and spatial resolution

Figure 4.2.6 shows that **most participants are interested in multiple years and long term statistics**, although a significant fraction of the participants are also interested in less-than-a-year time-series (24%) and single event data (43.5%).




Figure 4.2.6 Period of time participants are interested in

For what regard the spatial resolution at which the participants expect to use the data (Figure 4.2.7) it clearly appears that **high resolution (<10km) sea state data are of greater interest**, compared to low resolution and single point measurements.




Figure 4.2.7 Spatial resolution at which participants expect to use the data

4.2.5 Interest in sea state data

Figure 4.2.8 reveals that **the major interest for satellite wave data concerns the study of extreme events, the validation of wave models and the study of coastal processes**.

These three topics are followed by the study of *wave climate variability*, the study of *wave-current interactions*, the study of *air-sea interactions*, the study of *wave-ice interactions*, *data assimilation* and applications for *Marine Renewable Energy*. Other field of interest were also proposed (not shown), such as: *statistical downscaling*, *machine learning*, *ship hydrodynamics*, *offshore engineering*, and *seismic ambient noise characterization*.




Figure 4.2.8 Answers to the question “your interest in satellite wave data concerns?”

4.2.6 Interest in other Essential Climate Variables

To the question “have you used or are you planning to use **other CCI variables?**”, 101 participants answered **Sea Level**, 33 participants answered **Ocean Color**, 54 participants answered **Salinity**, 69 participants answered **Sea Surface Temperature**, and 42 participants answered **Sea Ice**.

4.2.7 Participation to the User Consultation Meeting and training session

To the question “do you plan to attend the User Consultation Meeting (UCM) in Brest, France (October 8th to 10th, 2019)”, 35% of the participants answered “Definitely yes” or “Probably” (Figure 4.3.6). In addition, many participants are interested in training sessions on *combining model and satellite data* (62%), *SAR data* (58%), *coastal altimetry* (56%), and *CFOSAT data* (29.5%). Hence, including a training session on one or several of the above-mentioned subjects to the User Consultation Meeting will likely attract more participants.

What next: do you plan to attend the User Consultation Meeting (UCM) in Brest, France (October 8th to 10th, 2019)?

180 responses




Figure 4.2.9 Interest of the participants in the User Consultation Meeting to be held in Brest in October 2019.

4.3 Second user survey

A second user survey was conducted in the context of the 2nd Sea State CCI User Consultation Meeting, which took place on 23-25 March 2021. The survey was open from 23 March to 17 June 2021. This survey was developed on LimeSurvey (<https://esa-survey.limequery.org/997962?lang=en>) with support from ESA communication manager P. Fischer and was broadcasted on the Sea State CCI website (<https://climate.esa.int/en/projects/sea-state/news/sea-state-cci-survey/>) and UCM mailing lists.

4.3.1 User profile and study area

As of June 17, a **total of 92 participants** had replied to the questionnaire. **25 countries** are represented. The most represented countries are France (16%), UK (11%), Germany (7%), Spain (4%), US (3%), and Australia (3%). The other represented countries are Brazil, Canada, Chile, China, Colombia, Costa Rica, Ireland, Italy, Japan, Mexico, Netherlands, New Zealand, Norway, Portugal, Romania, Russia, South Korea, Sweden. Similar to the first survey, we see on Fig. 4.3.1.1 that **the participants are mostly working in academic and research institutions (52%)**. The **main fields of interest are Physical Oceanography (52%), Climate (46%) and Spatial Oceanography (43%)**, followed by Coastal Engineering (25%), Marine Renewable Energy (23%) and Ocean Engineering (21%) (Fig. 4.3.1.2).




Figure 4.3.1.1 Answers to the question “What is your professional affiliation?”




Figure 4.3.1.2 Answers to the question “What are your disciplines of interest?”

66% were interested in the coastal region and were either working or not yet working on it. For the majority of the participants, **the 0-10km region is the coastal strip of most importance** (Fig. 4.3.1.3)




Figure 4.3.1.3 Answers to the questions “Are you interested in the coastal zone?” (left) and “What is the coastal strip of most importance?” (right)

Regarding the polar region, 46% are interested and either working or not yet working on this region. 27% did not show interest in the polar region.

4.3.2 Use of sea state data

According to this survey, the **main use of sea state observations is to validate numerical models** (55%), followed by **climate studies** (42%) and **extreme value analysis** (42%) (Fig. 4.3.2.1). **The most required temporal resolution is hourly** (63%), followed by long-term (27%) and monthly (27%) timescales (Fig. 4.3.2.2).




Figure 4.3.2.1 Answers to the questions “How do you make use of sea state data?”




Figure 4.3.2.2 Answers to the questions “What temporal resolution is required for your applications?”

In order to assess **which sea state parameters are the most important** to the user community, the participants were asked to rank 11 parameters (Significant wave height, Peak direction, Mean wave direction, Peak period, Mean period, Swell partitions, 1D spectrum, Full 2D wave spectrum, Mean square slope, Microseism sources, Wind) in terms of importance. The **significant wave height** was ranked first by 48% of the participants. **Then the wind** was ranked first by 10% and second by 21%. The **full 2D wave spectrum** was ranked first by 10%, and second by 5%. The mean wave direction was ranked second by 9% and third by 13%, while the peak period was ranked second by 10% and third by 11%.

To the questions “Which type of observations are you most familiar with?”, 32 participants ranked wave buoy first (67%), followed by satellite altimeter (17%) and ADCP (5%).

Participants were also asked to rank 9 sea state dataset that they ever used (Sea State CCI data, GLOBWAVE products, CMEMS in situ products, CMEMS model products, CMEMS satellite products, RADS, ECMWF reanalysis, NOAA reanalysis, IFREMER model hindcast). 20% ranked ECMWF reanalysis first, followed by Sea State CCI data (13%), CMEMS products, GLOBWAVE products, IFREMER reanalysis and NOAA reanalysis.

4.3.3 Experience with satellite data

Among the 92 participants, **14% had never used satellite sea state data**. Sea state satellite information was **considered as critical for the activity of 30%** of the participants.

The type of altimeter data that is **most used by the participants is the Level-2 along-track data, followed by simplified L3 products**. The time-delay product most used or needed is Re-analysis.




Figure 4.3.3.1 Answers to the questions “What type of satellite altimetry product do you mainly use?” (left) and “Which time-delay product do you use/need?” (right)

The participants were asked “What is currently the main limitation of satellite observation for your activity?”. For each of the seven proposed criteria (Spatial sampling, Temporal sampling, Accuracy, Time coverage, Available parameters, Accessibility of products, Ease of use), they were asked to select one of the free following answers: Not limiting, Partly limiting, Strongly limiting. The results are shown on Fig.4.3.3.2. **Temporal and spatial resolutions are the parameters considered as strongly limiting by the highest number of participants (>20%)**, followed by time coverage (13%). Accuracy and parameter availability are considered as partly limiting by more than 30%. Finally, **accessibility and ease of use are considered as not limiting by more than 20%**.

Moreover, a large number of the participants acknowledged that **temporal gaps (46%) and sampling discontinuity (40%)** affected the quality of their applications.

4.3.4 Experience with Sea State CCI products

Among the 92 participants, **37 (40%) had never used the sea state CCI products**. They were asked to select a reason (Fig. 4.3.4.1). **15% did not know where to find the data, 14% did not need this data, and 7% did not know how to use the data**. The other participants found the sampling or the accuracy too low for their application, or provided a reason not specified in the choices (see Appendix 1)




Figure 4.3.3.2 Answers to the questions “What is currently the main limitation of satellite observation for your activity?”




Figure 4.3.3.3 Answers to the questions “What is the main reason that explains why you haven’t used Sea State CCI data so far?”

The participants that already used the Sea State CCI products (20%) were asked which products they used among the L2, L3 and L4 products. **The L3 daily merged along-track product was the most used product** followed by L4 monthly gridded product and the L2 multi-mission along-track product (Fig 4.3.3.4). Moreover, **the most used significant wave height parameter is the swh denoised parameter**.




Figure 4.3.3.4 Answers to the questions “Which Sea State CCI product have you used?” (left) and “Which significant wave height parameter do you mainly use for your work?” (right)

Participants were asked if uncertainty information on sea state measurements is important for their activity and 40% answered positively while 14% answered negatively (Fig. 4.3.3.5). Among the proposed choices for uncertainty parameter:

1. High-resolution (20Hz) root-mean-square deviation from the mean (1Hz resolution)
2. Overall error statistics (bias, RMSE, SI) estimated from satellite-in situ platform comparison (mission specific)
3. Hs-dependent error model estimated from satellite-in situ platform comparison and applied to 1Hz data
4. Random error variance estimated from triple collocation analysis (e.g. altimeter, model, *in situ*)

Parameter #2 was ranked first by 35%, followed by parameter #1 (9%), parameter #4 (7%) and parameter #3 (5%).




Figure 4.3.3.5 Answers to the questions “Is uncertainty information on sea state parameters important to your activity ?”

Participants were asked to rank by order of priority which one of the following features they would you like to see in future Sea State CCI products:

- Extended time period with historical missions, from 1992 to 2002 (e.g. ERS, TOPEX...)
- Increased sampling density with more recent missions (e.g. CFOSAT, HY2A...)
- Inter-calibrated wind parameter
- Spectral swell parameter (direction, period) from SAR missions
- Spectral swell and wind-sea parameter (direction, period) from CFOSAT
- High-resolution wave heights in the coastal zone
- Increased spatial resolution of gridded product (currently 1°)
- Improved absolute and inter-mission calibration

- Waves in ice spectra
- Hs from Sentinel 1 SAR
- Wave period from SWH & sigma0 (either directly or via mss)

Figure 4.3.3.6 indicates that “**High-resolution wave heights in the coastal zone**” is the feature that was ranked first by the majority of the participants, followed by “Extended time period...” and “spectral swell parameter from SAR missions”.





Figure 4.3.3.6 Answers to the questions “which one of the following features would you like to see in future Sea State CCI products?” (ranked first)

Finally, the participants were asked to select the other CCI variable they plan to combine with Sea State CCI data in future analysis. The **first choice is “Sea Level” (22%), followed by Sea Ice (20%) and Sea Surface Temperature (16%)**. The other selected variables are: Ocean Colour, Sea Surface Salinity; Greenhouse gases, Ice Sheets (Antartica and Greenland), Aerosol, Biomass, and Land Cover.



5. Consolidated list of User Requirements

Based on the above material, we can highlight the relevant aspects of the user requirements for the CCI Sea State project. The specific data requirements as defined in the ESA Statement of Work are recalled in section 5.2, and are linked to the user requirements when appropriate.

5.1 Top level requirements

- **resolution:** There is a clear need for data at a resolution finer than the 25 km mentioned in GCOS, in particular in the coastal regions (here defined as the combination of depths under 100 m and distances to shore under 300 km). Most of the surveyed users would like to have **10 km or less**. Given the resolution of standard satellite processing, this is really calling for improved tracking and/or denoising algorithms, and by itself it justifies the effort made on the Sea State CCI.
- **coverage in space and time:** Given that a large fraction of users are interested into **single events** (80 out of 184 participants in the first survey), or considered that **spatial and temporal sampling are strongly or partly limiting the use of the data** (~40 out of 92 participants in the second survey), this clearly highlights the sampling issue of satellite data sets. Most events are missed, except through their associated swell fields. There is thus a need that will not be fully addressed in the Phase 1 of Sea State CCI for combining wave models and data or expanding on level 3 and 4 products such as fireworks, storm catalogs (associated with storm tracks ...). Regarding ice-covered regions, 49% of the participants of the second survey indicated they were interested in the polar region.
- **Stability:** not surprisingly most users identified here are interested into long-term statistics of sea state variable, with or without a climate change aspect. Given that

many users (101/184 in survey 1 and 20/92 in survey 2) mentioned their intent to combine sea level with sea state, it is logical to reframe the requirements on sea state stability in terms of total sea level (e.g. Dodet et al., 2019, see also [15] , Marcos et al. 2019). The **need** on wave height trend accuracy for mean values and extremes is thus **under 1 mm/year for coastal areas**, with a similar need for wave periods that should be quantified. It is not at all clear that such a low value can be achieved with today's spatial coverage, and how bringing models forced by winds with dubious trends can be used for this. At any rate, even the GCOS requirement of 5 cm/decade, when achieved, should be enough to confirm or disprove the 0.5% per year (around 1 cm per year) trend associated with wave power trends up to 2.5% per year reported by Reguero et al. (2019).

- **Sea State Variables:** It is not just the **wave heights**. For many reasons (energy flux, extreme sea levels...) the **full directional spectra**, or the **periods** and **directions** have a very important role, and this is well recognized in the user survey. It is thus very important to use both altimeters (via the cross-section) and wave-resolving instruments (SARs, SWIM on CFOSAT) for constraining the sea state climate. It may be surprising that fewer users would like to use partition data, but this may be due to the fact that little such data is available and the definition can be a bit fuzzy and method-dependent (e.g. Portilla et al. 2009). We will thus engage the user community (at UCM and through training events) on this question and see how the usefulness of such data can be improved. Given the very few users of SAR data, making the data more accurate and also more user-friendly is a key aspect.

5.2 Specific data requirements

ID	Requirement description	Source	Comment
DR-001	In Phase 1 the Sea_state_cci project shall develop an initial 18-year data set (2002- 2020), and shall provide, as a minimum, L2 products and higher level merged product time-series that shall collectively include the following sea state variables: Significant wave height; Directional wave spectrum; Mean wave period; Peak wave period; Mean wave direction at the peak of the spectrum; Appropriate derived-variables and supporting variables; Other variables required by the Climate Science Community.	ESA TR-2, TR-15, TR-16	
DR-002	Each CCI project team (the contractor) shall integrate data from the Copernicus Sentinels and other key satellite missions within the relevant CCI processing systems and ECV data products.	ESA R-16	Sentinel 1, 3 & 6 included in baseline Sentinel 2 in option 10.
DR-003	Each CCI project team (the contractor) shall ensure that the system is adequately dimensioned to accommodate the growing volumes of input and output data, and the increasing computational loads needed to process, reprocess, quality control, validate, and disseminate multi-decadal, global, ECV	ESA R-17	

	data products, of the required climate quality, in a timely manner.		
DR-004	Sea_state_cci shall directly address GCOS Action O33.	ESA TR-1	
DR-005	The Sea_state_cci project shall develop and deliver Sea State ECV products primarily derived from satellite measurements.	ESA TR-7	
DR-006	The Sea_state_cci project shall deliver validated prototype products using agreed validation methods and metrics developed within a research environment to climate science users for assessment and feedback.	ESA TR-8	
DR-007	Sea_state_cci products shall cover the global ocean, including full coverage of both northern and southern hemispheres as far as possible.	ESA TR-9	
DR-008	All Sea_state_cci products shall cover the full mission lifetimes of the satellite missions selected.	ESA TR-10	
DR-009	Sea_state_cci products shall be available to users as Level-1 (where appropriate), Level-2 and Level-3 product versions, and potentially as higher-level derived products if required by the users.	ESA TR-11	
DR-010	Sea_state_cci products shall include aggregated versions of the data as required by climate science users (eg. daily, monthly, seasonally and annually).	ESA TR-12	
DR-011	Digital Object Identifiers (DOI) shall be assigned to all ECV data sets made publicly available.	ESA TR-13	
DR-012	As part of data merging methods, time-dependent and sampling biases in products from different instruments shall be investigated, and strategies shall be developed and implemented to correct for these effects.	ESA TR-17	WP2*70
DR-013	A common set of auxiliary/supporting data shall be developed and used for all satellite missions used within the Sea_state_cci project.	ESA TR-20	WP4200
DR-014	The Sea_state_cci project shall explore techniques using satellite measurements made at different frequencies (eg. C, S, Ku, Ka bands) to address GCOS Sea State ECV requirements.	ESA TR-22	
DR-015	The Sea_state_cci project shall develop innovative merging strategies and tools for sea state products generation.	ESA TR-23	WP2*70
DR-016	The Sea_state_cci project shall provide a validated estimate of uncertainty for each data product at product grid/pixel level using the approach of [ESA RD-33].Uncertainties shall be reported within the ECV products for every geophysical measurement.	ESA TR-25, TR-39, TR-40	
DR-017	The method used to derive and validate uncertainties, the characteristics of those uncertainty estimates and advice on how uncertainty estimates are to be used for each product shall	ESA TR-26	

	be fully reported in the PUG.		
DR-018	User requirements for ECV product uncertainties shall be included in the user requirements analysis, including how the uncertainties should be expressed and used in the Sea_state_cci ECV products (e.g. how should the uncertainties be broken down into their random and systematic components).	ESA TR-30	WP1000
DR-019	The Contractor shall conduct significant research and development and explore innovative approaches and algorithms that could address known weaknesses in sea state retrievals from satellite data sets.	ESA TR-31	WP2000
DR-020	The Contractor shall conduct research and development and explore new algorithms to address crossing seas.	ESA TR-33	
DR-021	The Contractor shall investigate and account for satellite instrument biases, particularly regarding earlier less well calibrated instruments taking account of changes in calibration with instrument aging.	ESA TR-35	
DR-022	Based on the outcome of the Round Robin, The contractor shall select a set of definitive retrieval algorithms to be applied to data from different instruments.	ESA TR-36, TR-37, TR-38	WP2100
DR-023	The Contractor shall ensure the new capabilities of Copernicus Sentinel-3 SRAL (and in future Copernicus Sentinel-6) SAR altimeter instruments are fully exploited for the retrieval of sea state ECV products.	ESA TR-41	
DR-024	The Contractor shall ensure the new capabilities of Copernicus Sentinel-1 SAR imager instruments are fully exploited for the retrieval of the sea state ECV products.	ESA TR-42	
DR-025	A full validation of all sea state ECV products produced shall be performed against metrics pre-defined defined by the contractor and endorsed by the user community.	ESA TR-43, TR-44	WP4500
DR-026	Validation shall quantify the uncertainty of the sea state ECV products as well as the quality of the product uncertainty estimates themselves.	ESA TR-45	WP4500
DR-027	The long-term stability of all ECV time series delivered shall be assessed.	ESA TR-46	WP4500
DR-028	A database of relevant and ideally independent <i>in situ</i> Fiducial Reference Measurements and satellite measurements (ISDB) shall be developed to serve the Sea_state_cci project validation, research and development needs.	ESA TR-47	WP4100
DR-029	All measurements in the ISDB shall include uncertainty estimates. The methods used to derive and validate ISDB uncertainties and the characteristics of those uncertainty estimates for each product shall be fully reported in the PUG.	ESA TR-48, TR-50	
DR-030	The suite of Sea_state_cci ECV products produced shall be	ESA	

	made publicly available together with the validation results immediately following the completed validation.	TR-54	
--	--	-------	--

6. References

- [1] EUMETSAT Position Paper on Observation Requirements for Nowcasting & Very Short Range Forecasting in 2015-2025, B.W. Golding, S. Senesi, K. Browning, B. Bizzarri, W. Benesch, D. Rosenfeld, V. Levizzani, H. Roesli, U. Platt, T. E. Nordeng, J. T. Carmona, P. Ambrosetti, P. Pagano, and M. Kurz
- [2] EUMETSAT Position Paper - Requirements of Observation for Regional NWP, N. Gustafsson, M. Capaldo, B. O. Estrada, and J. Quiby
- [3] EUMETSAT Position Paper - Requirements of Observation for Global NWP, J. Eyre, J.-N. Thépaut, J. Joiner, L. P. Riishojgaard and F. Gérard
- [4] An Ocean Theme for the IGOS Partnership, Final Report from the Ocean Theme Team, IGOS, 36 pp, www.igospartners.org
- [5] IOC/UNESCO Report on “Observing the Ocean in the 21st Century”, ed. by Koblinsky, C.J. and Smith, R.N., 2001, 604 pp.
- [6] Global Observing Systems Information Center, Space-Based Data Requirements for GOOS, http://www.gosic.org/space/GOOS_space.htm
- [7] GMES Sentinel 3 Mission Requirements Document, ESA document EOP-SMO/1151/MD-md issue 1.0, 2004.
- [8] NPOESS Integrated Operations Requirements Document, 2001.
- [9] EUMETSAT Satellite Application Facility on Ocean and Sea Ice, User Requirements Document, EUMETSAT document number SAF/OSI/M-F/MGT/RQ/011, Version 2.0, 2005.
- [10] GCOS, 2006. SYSTEMATIC OBSERVATION REQUIREMENTS FOR SATELLITE-BASED DATA PRODUCTS FOR CLIMATE. Supplemental details to the satellite-based component of the “Implementation Plan for the Global Observing System for Climate in Support of the UNFCCC (GCOS – 92)”.
- [11] (WMO/TD No. 1143), The Second Report on the Adequacy of the Global Observing Systems for Climate in Support of the UNFCCC, 2003. See <http://www.wmo.ch/web/gcos/gcoshome.html>
- [12] GCOS, 2011. SYSTEMATIC OBSERVATION REQUIREMENTS FOR SATELLITE-BASED DATA PRODUCTS FOR CLIMATE, 2011 Update. Supplemental details to the satellite-based component of the “Implementation Plan for the Global Observing System for Climate in Support of the UNFCCC (2010 Update)” December 2011, **GCOS – 154**.
- [13] COST Action 714 Working Group 3, W. G., Measuring and analysing the directional spectra of ocean waves. Office for Official Publications of the European Communities, Luxembourg, 2005. ISBN 92-898-0003-8.
- [14] Ardhuin, F., Ocean waves in geosciences. 2019. doi:10.13140/RG.2.2.16019.78888/5.

[15] [Ardhuin, F., J. E. Stopa](#), B. Chapron, F. Collard, [R. E. Jensen, J. Johannessen](#), R. Husson, A. Mouche, [M. Passaro, G. Quartly, V. Swail and I. Young](#). Observing sea states, *Frontiers in Marine Science* (in review https://www.researchgate.net/publication/328791679_Observing_sea_states).

Allen, P. A. and Hoffman, P. F., “Extreme winds and waves in the aftermath of a neoproterozoic glaciation,” *Nature*, 433, 123–127, 2005. See commentary by D. J. Jerolmack and D. Mohrig, *Nature*, 2005, vol. 436, p. E1.

Ardhuin, F., Drake, T. G., and Herbers, T. H. C., “Observations of wave-generated vortex ripples on the North Carolina continental shelf,” *J. Geophys. Res.*, 107, C10, 2002. doi:10.1029/2001JC000986.

Ashton, A., Murray, A. B., and Arnault, O., “Formation of coastline features by large-scale instabilities induced by high-angle waves,” *Nature*, 414, 296–299, 2001. doi:10.1038/35104541.

Aucan, J., Hoeke, R., Merrifield, M.A., 2012. Wave-driven sea level anomalies at the Mid-way tide gauge as an index of North Pacific storminess over the past 60 years. *Geophys. Res. Lett.* 39, 1–6. <http://dx.doi.org/10.1029/2012GL052993> .

Autret, R., Dodet, G., Fichaut, B., Suanez, S., David, L., Leckler, F., Ardhuin, F., Ammann, J., Grandjean, P., Lallemand, P., and Filipot, J.-F., “A comprehensive hydro-geomorphic study of cliff-top storm deposits on Banneg island during winter 2013-2014,” *Marine Geology*, 382, 37–55, 2016. doi:10.1016/j.margeo.2016.09.014.

Bernard, P., “Historical sketch of microseisms from past to future,” *Phys. Earth Planetary Interiors*, 63, 145–150, 1990. doi:0031-9201(90)90013-N.

Bouchette, F., Schuster, M., Ghienne, J.-F., Denamiel, C., Roquin, C., Moussa, A., Marsaleix, P., and Durringer, P., “Hydrodynamics in holocene lake mega-chad,” *Quaternary Research*, 73, 226–236, 2010. doi:10.1016/j.yqres.2009.10.010.

Cox, R., “Very large boulders were moved by storm waves on the west coast of Ireland in winter 2013–2014,” *Marine Geology*, p. in press, 2018. doi:10.1016/j.margeo.2018.07.016.

de Leeuw, G., E. L. Andreas, M. D. Anguelova, C. W. Fairall, E. R. Lewis, C. O’Dowd, M. Schulz, and S. E. Schwartz (2011), Production flux of sea spray aerosol, *Rev. Geophys.*, 49, RG2001, doi:10.1029/2010RG000349.

Doble, M. J. (2003). *Pancake ice formation in the Weddell Sea. Journal of Geophysical Research*, 108(C7). doi:10.1029/2002jc001373

Dodet, G., Leckler, F., Sous, D., Ardhuin, F., Filipot, J. F., & Suanez, S. (2018). *Wave Runup Over Steep Rocky Cliffs. Journal of Geophysical Research: Oceans*. doi:10.1029/2018jc013967

Donelan, M. A., “On the decrease of the oceanic drag coefficient in high winds,” *J. Geophys. Res.*, 123, C2, 1485–1501, 2015. doi:10.1002/2017JC013394.

Jähne, B. and Haussecker, H., “Air-water gas exchange,” *Annu. Rev. Fluid Mech.*, 30, 443–469, 1998.

Gerling, T. W., “Partitioning sequences and arrays of directional ocean wave spectra into component wave systems,” *J. Atmos. Ocean Technol.*, 9, 444–458, 1992.

Grevemeyer, I., Herber, R., and Essen, H.-H., “Microseismological evidence for a changing wave climate in the northeast Atlantic Ocean,” *Nature*, 408, 349–351, 2000.

Hansen, J., Sato, M., Kharecha, P., & von Schuckmann, K. (2011). *Earth’s energy imbalance and implications. Atmospheric Chemistry and Physics*, 11(24), 13421–13449. doi:10.5194/acp-11-13421-2011

Hansen, J., Sato, M., Hearty, P., Ruedy, R., Kelley, M., Masson-Delmotte, V., Russell, G., Tselioudis, G., Cao, J., Rignot, E., Velicogna, I., Tormey, B., Donovan, B., Kandiano, E., von Schuckmann, K., Kharecha, P., Legrande, A. N., Bauer, M., and Lo, K.-W., “Ice melt, sea level rise and superstorms: evidence from paleoclimate data, climate modeling, and modern observations that 2°C global warming could be dangerous,” *Atmos. Chem. Phys.*, 16, 3761–3812, 2016. doi:10.5194/acp-16-3761-2016.

Hanson, J. L. and Phillips, O. M., “Automated analysis of ocean surface directional wave spectra,” *J. Atmos. Ocean Technol.*, 18, 277–293, 2001.

Kennedy, A. B., Mori, N., Zhang, Y., Yasuda, T., Chen, S.-E., Tajima, Y., Pecor, W., and Toride, K., “Observations and modeling of coastal boulder transport and loading during super typhoon haiyan,” *Coastal Eng. Japan*, 58, p. 1640004, 2016. doi:10.1142/S0578563416400040.

Lygre, A., and H. E. Krogstad, 1986: Maximum entropy estimation of the directional distribution in ocean wave spectra. *J. Phys. Oceanogr.*, 16, 2052–2060.

Marcos, M., Wöppelmann, G., Matthews, A., Ponte, R. M., Birol, F., Arduin, F., ... Stopa, J. E. (2019). Coastal Sea Level and Related Fields from Existing Observing Systems. *Surveys in Geophysics*. doi:10.1007/s10712-019-09513-3

Passaro, M., Z. A. Nadzir, G. D. Quartly, Improving the precision of sea level data from satellite altimetry with high-frequency and regional sea state bias corrections, *Remote Sensing of Environment* 218 (2018) 245–254, doi: /10.1016/j.rse.2018.09.007

Poate, T. G., McCall, R. T., and Masselink, G., “A new parameterisation for runup on gravel beaches,” *Coastal Eng.*, 117, 176–190, 2016.

Reguero, B. G., Losada, I. J., & Méndez, F. J. (2019). *A recent increase in global wave power as a consequence of oceanic warming. Nature Communications*, 10(1). doi:10.1038/s41467-018-08066-0

Reul, N. and Chapron, B., "A model of sea-foam thickness distribution for passive microwave remote sensing applications," *J. Geophys. Res.*, 108, C10, p. 3321, 2003. doi:10.1029/2003JC001887.

Semedo, A., Sušelj, K., Rutgersson, A., Sterl, A., 2010. A Global View on the Wind Sea and Swell Climate and Variability from ERA-40. *Journal of Climate* 24, 1461–1479. <https://doi.org/10.1175/2010JCLI3718.1>

Shapiro, N. M., Campillo, M., Stehly, L., and Ritzwoller, M. H., "High-resolution surface-wave tomography from ambient seismic noise," *Science*, 307, 1615–1617, 2005. doi:10.1111/j.1365-246X.2006.03240.x.

Shimura, T., Mori, N., and Hemer, M. A., "Variability and future decreases in winter wave heights in the western north pacific," *Geophys. Res. Lett.*, 43, 2716–2722, 2016. doi:10.1002/2016GL067924.

Stopa, J. E., Ardhuin, F., and Girard-Ardhuin, F., "Wave climate in the arctic 1992-2014: seasonality and trends," *The Cryosphere*, 10, 1605–1629, 2016. doi:10.5194/tc-10-1605-2016.

Squire, V., Dugan, J., Wadhams, P., Rottier, P., and Liu, A., "Of ocean waves and sea ice," *Annu. Rev. Fluid Mech.*, 27, 3, 115–168, 1995.

Stopa, J., P. Sutherland, F. Ardhuin, Strong and highly variable push of ocean waves on Southern Ocean sea ice, *Proc. Nat. Acad. Sci.*, 115(23), 5861-5865, doi :10.1073/pnas.1802011115

Sutherland, P., D. Dumont, Marginal ice zone thickness and extent due to wave radiation stress, 2018, *J. Phys. Oceanogr.*, doi : 10.1175/JPO-D-17-0167.1

Thomson, J., S. Ackley, F. Girard-Ardhuin, F. Ardhuin, A. Babanin, G. Boutin, J. Brozena, S. Cheng, C. Collins, M. Doble, C. Fairall, P. Guest, C. Gebhardt, J. Gemmrich, H. C. Graber, B. Holt, S. Lehner, B. Lund, M. H. Meylan, T. Maksym, F. Montiel, W. Perrie, O. Persson, L. Rainville, W. E. Rogers, H. Shen, H. Shen, V. Squire, S. Stammerjohn, J. Stopa, M. M. Smith, P. Sutherland, P. Wadhams, Overview of the Arctic Sea State and Boundary Layer Physics Program, *J. Geophys. Res.*, 2018. doi :10.1002/2018JC013766

Tran, N., Vandemark, D., Labroue, S., Feng, H., Chapron, B., Tolman, H. L., Lambin, J., and Picot, N., "The sea state bias in altimeter sea level estimates determined by combining wave model and satellite data," *J. Geophys. Res.*, 115, p. C03020, 2010. doi:10.1029/2009JC005534.

Veron, F., "Sea spray," *Annu. Rev. Fluid Mech.*, 47, 507–538, 2015. doi:10.1146/annurev-fluid-010814-014651.

Wiese, A., Staneva, J., Schulz-Stellenfleth, J., Behrens, A., Fenoglio-Marc, L., and Bidlot, J.-R.: Synergy of wind wave model simulations and satellite observations during extreme events, *Ocean Sci.*, 14, 1503-1521, <https://doi.org/10.5194/os-14-1503-2018>, 2018.

Young, I. R., & Donelan, M. A. (2018). *On the determination of global ocean wind and wave climate from satellite observations*. *Remote Sensing of Environment*, 215, 228–241. doi:10.1016/j.rse.2018.06.006

Appendix 1 - Open comments from survey participants

First survey

- *Right now I am just a peripheral user of satellite data. However me and my group are planning to use this data more, as we would like to widen our wave observations away from single buoy / point observations.*
- *Thanks for the work done*
- *Keen to try your product soon*
- *At this stage, I'm still trying to process the satellite data that I downloaded on the CMEMS server (Jason 3 along track significant wave height) and once I'm done with that, I will be to comment and make suggestions when needed.*
- *I've never used satellite measurements but would be interested in possible applications in the nearshore*
- *The satellite data are not open data, their availability depends on the country of satellite ownership, the country of user's residency, the nature of the organizations of all the interested parties and more.*
- *hope it works well for u!*
- *The datasets produced by the CCI might last for a decade so it is important to make the most of the project and ensure all parties (users, scientists, engineers, industry, ...) are satisfied as best as possible.*
- *I feel rather out of touch, due to other work commitments, but very happy to see all of this happening and I hope to be more involved later*
- *Nice survey.*
- *thanks for collecting this information*
- *thanks*
- *Thank you so much for the initiative. I really appreciate it!*
- *Thank you for the great work you do.*
- *None*
- *Thanks for the great work you are doing in this interdisciplinary area. Two further comments in support of your excellent work in this area:
1 - if an interdisciplinary user like me / my group members / isn't using a product it might be because they didn't know about it yet.
2 – with data-driven (e.g. machine learning) approaches, we just need lots of data and not to worry too much about each individual type of observable. Sometimes it is better not to pre-empt what is a useful observable.*
- *It would be great to have a page with a list of data sources for all available historical and NRT SAR L2, L3 datasets including ERS-1/2, Envisat, and Sentinel-1A/B together with their pros and cons (e.g. good quality/bad quality/no quality), similarly for buoy data (both directional and otherwise) and also for altimetry data.*

- *Just a single summary page of links and pros and cons of using the data.*
- *I would prefer to have access to raw (less smooth) data, and do my own processing to them.*
- *It will have a great deal if one can be able to validate the wave models with the satellite data*
- *I'm not sure about merged altimeter data provided by IFREMER. But I contacted the concerned person I didn't receive any response. Then it's like use with your own risk. Can someone help me with my doubts in data inhomogeneity issues?*

Second user survey

19. If you haven't used the CCI products yet, what is the main reason? [Comment]

- *But going to use them soon*
- *Don't know if it's useful for this application*
- *Hadn't thought about it*
- *Have not decided the best way to use it yet*
- *I already used Globwave data and CMEMS satellite data*
- *I did not know about the dataset*
- *I didn't have the chance yet but I am planning to use.*
- *I didn't know about the existence of this data*
- *I may want this data as the project progresses to a later stage*
- *I was not aware about the existence of this data set*
- *I will probably use it in the next future*
- *just find out that they are exist*
- *Not had time yet!*
- *Operational mission with limited time, however, future activities we will consider.*
- *other set available*
- *Unaware until recently*

27. What sort of statistics would you like to find in a higher level product (gridded summary products)? [Comment]

- *Benjamin & Feir index, cross seas index*
- *Bias, R, RMSE, SI*
- *Monthly max, seasonal max*
- *Will derive from time series*

28b You have indicated that you plan to combine Sea State CCI data with other CCI variables. Would you like to see improved inter-operability between these datasets?

[Comment]

- *Consistency of sea ice / open water classification. Potential wave propagation and dampening in marginal sea ice zones.*
- *Improving the precision of sea level data from satellite altimetry with high-frequency and regional sea state bias corrections*

29. Do you see any information that is missing from the metadata and that you would like to see in future versions?

- *The present "Sea state CCI Product User Guide, V1.0" has a comprehensive list of variables for all datasets (L2P, L3, L4). However, a clear definition of all variables is*

not given, for example, what is the meaning (in L4) of a monthly value for "swh_mean: mean of median significant wave height values"? (In this example, what and how is being averaged?, I guess one averages several values of median_swh, but is it a spatial average or a temporal average). So, my suggestion goes to include in an Annex a clear mathematical/physical definition of all variables.

30. Do you have any other suggestions?

- *I downloaded the data rapidly with no difficulties, keep on!*
- *I would suggest to make the CCI data sets available through CMEMS and other major metocean portals. This could allow to increase the dissemination. An coordination with CMEMS to better highlight the complementarities between ESA CCI and CMEMS sea state satellite product would be also useful.*
- *More frequent data releases, even if the data are experimental only (to keep up with rapid developments in other projects and countries)*
- *Provide the inter-satellite cross-over database (for altimeter vs.SAR, altimeter vs. wave scatterometer and SAR vs. wave scatterometer) for further analysis of possible synergistic new sea state products*
- *Since the "swh_denoised" is the swh best estimate from the data, I suggest to provide solely this variable, as providing the others can be confusing and misleading.*
- *Some questions in the survey seemed to ignore the existence of level-4 products, which are the only ones I have used so far. For instance, one question only had L1, L2 and L3 as possible answers. The next one I think was asking whether we use raw or denoised/corrected SWH, but this is not relevant for L4 data, which, as far as I understand, only provide denoised/corrected SWH.*
- *Question 28. has an issue: I haven't found boxes for Yes/No answer, but then 28b assumes that I ticked Yes (28a nowhere in sight). Actually, my answer to this question is No. So I had no other choice than not providing answers to either 28 or 28b.*
- *Thank you for your work!*